

**On a generalization of the Fay-Sato identity
for KP Baker functions and
its application to constrained hierarchies**

LEONID DICKY

Math. Department, University of Oklahoma
Norman, OK 73019, USA
e-mail: ldickey@uoknor.edu
and

WALTER STRAMPP

Fachbereich 17-Mathematik/Informatik
Universität-GH Kassel
Holländische Strasse 36
34109 Kassel, Germany
e-mail: strampp@hrz.uni-kassel.de

Abstract

Some new formulas for the KP hierarchy are derived from the differential Fay identity. They proved to be useful for the k -constrained hierarchies providing a series of determinant identities for them. A differential equation is introduced which is called “universal” since it plays an important role for all the k -constrained hierarchies. In the cases $k = 1, 2$ and 3 explicit formulas are presented, in all the others recurrence relations are given which enable one to obtain the identities.

0. Introduction. This paper is a result of our discussions about the meaning and the significance of a proposition in the article [1] (Proposition 2). We found out that a broader problem can be posed and solved using a more general language. Virtually, we suggest a new type of problems for the KP and, especially, k -constrained KP hierarchies. The explicit identities are obtained for the smallest values of k , $k = 1, 2$ and 3 ; for higher values we have recurrence formulas, i.e. a recipe to get new identities. Calculations become cumbersome. We hope that there exists a more general approach allowing one to circumvent these complicated calculations at all.

The so-called differential Fay identity for the KP hierarchy (proven by Sato for the most general τ functions) is well-known (see, e.g., [3]). It can be written in the following form using the Baker function $\hat{w}(t, z_1) \exp \sum t_i z_1^i$. Let $\mathbf{G}$ be a shift operator defined as

$$\mathbf{G}F(t_1, t_2, \dots) = F(t_1 + 1/z_2, t_2 + 1/(2z_2^2), t_3 + 1/(3z_2^3), \dots)$$

where z_2 is another parameter and let $\mathbf{G}_1 = \mathbf{G} - 1$. Then

$$(z_1 - z_2)\mathbf{G}_1\hat{w} = -\mathbf{G}_1\hat{w}' + \mathbf{G}_1w_1 \cdot \mathbf{G}_1\hat{w} + \mathbf{G}_1w_1 \cdot \hat{w} - \hat{w}'. \quad (0.1)$$

Here and further $\hat{w}$ is always $\hat{w}(t, z_1)$ and w_1 is the first non-trivial coefficient in the expansion $\hat{w} = \sum_0^\infty w_i z_1^{-i}$ (recall that $w_0 = 1$). Let

$$\mathbf{H} = -\partial + \mathbf{G}_1w_1 + \hat{w} \cdot \text{res}_{z_1} \quad (0.2)$$

be an operator that acts on series $\sum a_{ij} z_1^{-i} z_2^{-j}$. Eq.(0.1) can be represented as

$$(z_1 - z_2)n^{(1)} = \mathbf{H}(n^{(1)}) - \hat{w}', \quad \text{where } n^{(1)} = \mathbf{G}_1\hat{w}. \quad (0.3)$$

In fact, this identity is equivalent to the hierarchy itself since, expanding it in powers of z_1^{-1} and z_2^{-1} , all the $\partial_j w_i$ can be obtained.

Let us think on the last term of (0.3) as on $-\partial_1 \hat{w}$. We are going to generalize this identity by showing that one can construct quantities $n^{(k)} = \sum_1^\infty n_j^{(k)} z_2^{-j}$ where $n_j^{(k)} = P_j^{(k)} \hat{w}$ and $P_j^{(k)}$ are differential operators in ∂_i such that

$$(z_1 - z_2)n^{(k)} = \mathbf{H}(n^{(k)}) - \partial_k \hat{w}, \quad \partial_k = \partial/\partial t_k. \quad (0.4)$$

Explicit formulas for $n^{(k)}$ are obtained only for $k = 1, 2, 3$. In the case of larger k we give the recurrence formulas for $n_j^{(k)}$.

These identities have an interesting application in the theory of constrained KP hierarchies. The k -constraint for the KP hierarchy is $L^k = L_+^k + q\partial^{-1}r$. An important role for all the k -constraints plays an equation which we call “universal”. This is

$$f' + 1 - z_2 f + Qf - \mathbf{G}_1w_1 \cdot f = 0 \quad (0.5)$$

where Q is any function. The universal equation has the only solution in the form of a series in z_2^{-1} , $f_Q = \sum_1^\infty f_j z_2^{-j}$. The coefficients f_j are functions, one can find that $f_Q = z_2^{-1} + Qz_2^{-2} + \dots$. It will be shown that for the k -constrained hierarchy the quantities $n^{(k)}/\partial_k \hat{w}$ with the above $n^{(k)}$ satisfy the universal equation with $Q = q'/q$. Therefore $n^{(k)}/\partial_k \hat{w} = f_Q$ and they do not

depend on z_1 . The quantity $n^{(k)}$ happens to be a product of a series in z_1^{-1} and a series in z_2^{-1} . Hence, the relations $P_j^{(k)} w_i = f_j \cdot \partial_k w_i$, $j \geq k$ hold where $P_j^{(k)}$ are the above differential operators. This implies that the matrix $P_j^{(k)} w_i$ is of rank 1 and all of its second order minors vanish. This supplies one with infinitely many identities. The first of them, with $j_1 = 1$ and $j_2 = 2$ and arbitrary i_1 and i_2 , form a necessary and sufficient condition that a Baker function of the KP hierarchy satisfies the k th constraint.

1. The KP hierarchy. Let us recall some terminology.

Let

$$L = \partial + u_1 \partial^{-1} + u_2 \partial^{-2} + \dots$$

be a pseudodifferential operator, then

$$\partial_n L = [L_+^n, L], \text{ where } \partial_n = \partial / \partial t_n$$

are the equations of the KP hierarchy,

$$L = \hat{w}(t, \partial) \partial \hat{w}^{-1}(t, \partial), \quad \hat{w}(t, \partial) = \sum_0^\infty w_k(t) \partial^{-k}, \quad w_0 = 1,$$

and

$$\psi(t, z) = \hat{w}(t, z) \exp \xi = \hat{w}(t, \partial) \exp \xi \text{ where } \xi(t, z) = \sum_1^\infty t_k z^k$$

is the (formal) Baker function.

It is easy to see that the equations

$$\partial_n \hat{w}(t, \partial) = -L_-^n \hat{w}(t, \partial)$$

imply the above hierarchy equations for L . Then ψ satisfies the equations

$$L\psi = z\psi, \quad \partial_n \psi = L_+^n \psi.$$

The Schur polynomials are defined by

$$\exp \xi = \sum_0^\infty p_n(t) z^n, \quad t = (t_1, t_2, \dots).$$

Let $\tilde{\partial} = (\partial_1, \partial_2/2, \partial_3/3, \dots)$. Then $p_n(\tilde{\partial})$ are differential operators denoted by p_n^+ .

2. The τ -function and the differential Fay identity. The τ -function is defined by the equality

$$\hat{w}(t, z) = \frac{\tau(t - [z^{-1}])}{\tau(t)} \tag{2.1}$$

where $\tau(t) = \tau(t_1, t_2, \dots)$ and $\tau(t - [z^{-1}]) = \tau(t_1 - 1/z, t_2 - 1/(2z^2), t_3 - 1/(3z^3), \dots)$. The existence of such a function is not obvious and must be proven. An important step in the proof is establishment of a relation

$$\frac{\hat{w}(t - [z_2^{-1}], z_1)}{\hat{w}(t, z_1)} = \frac{\hat{w}(t - [z_1^{-1}], z_2)}{\hat{w}(t, z_2)} \tag{2.2}$$

which, in its turn, easily follows from (2.1). We need further a corollary of this equality that can be obtained expanding it in powers of z_1^{-1} :

$$\frac{\hat{w}'(t, z_2)}{\hat{w}(t, z_2)} = -w_1(t - [z_2^{-1}]) + w_1(t). \quad (2.3)$$

The relation (2.2) states that $\hat{w}(t - [z_2^{-1}], z_1)/\hat{w}(t, z_1)$ is symmetric with respect to z_1 and z_2 and $(z_2 - z_1)\hat{w}(t - [z_2^{-1}], z_1)/\hat{w}(t, z_1)$ is skew symmetric. We can do a more precise statement.

Lemma. *A relation*

$$(z_2 - z_1) \frac{\hat{w}(t - [z_2^{-1}], z_1)}{\hat{w}(t, z_1)} = -\frac{\hat{w}'(t, z_1)}{\hat{w}(t, z_1)} + \frac{\hat{w}'(t, z_2)}{\hat{w}(t, z_2)} + z_2 - z_1$$

holds.

Proof. As it is easy to check, this is nothing but the differential Fay identity (see, e.g., [3])

$$\begin{aligned} & \partial \tau(t - [z_1^{-1}]) \tau(t - [z_2^{-1}]) - \tau(t - [z_1^{-1}]) \partial \tau(t - [z_2^{-1}]) \\ &= (z_2 - z_1) \{ \tau(t - [z_1^{-1}]) \tau(t - [z_2^{-1}]) - \tau(t - [z_1^{-1}] - [z_2^{-1}]) \tau(t) \} \end{aligned}$$

expressed in terms of $\hat{w}(t, z)$ instead of $\tau(t)$. This identity follows from Sato's bilinear identity. $\square$

Shifting the argument $t \mapsto t + [z_2^{-1}]$ and taking into account (2.3) one gets

$$\begin{aligned} & (z_1 - z_2) \{ \hat{w}(t + [z_2]^{-1}, z_1) - \hat{w}(t, z_1) \} \\ &= -\hat{w}'(t + [z_2^{-1}], z_1) + \{ w_1(t + [z_2^{-1}]) - w_1(t) \} \hat{w}(t + [z_2^{-1}], z_1). \end{aligned}$$

For simplicity, let us introduce notations: $\hat{w}$ is always $\hat{w}(t, z_1)$, $\mathbf{G}$ will denote the shift $\mathbf{G}\hat{w} = \hat{w}(t + [z_2^{-1}], z_1)$. Thus, the statement of the lemma takes the form

$$(z_1 - z_2)(\mathbf{G}\hat{w} - \hat{w}) = -\mathbf{G}\hat{w}' + (\mathbf{G}w_1 - w_1)\mathbf{G}\hat{w}.$$

This equation contains all the equations of the hierarchy. They can be obtained by an expansion of the equation in powers of z_2^{-1} taking into account that $\mathbf{G} = \exp(\sum_1^\infty \partial_k z_2^{-k}/k) = \sum_0^\infty p_n^+ \cdot z_2^{-n}$:

$$z_1 \hat{w}' = p_2^+ \hat{w} - \hat{w}'' + w_1' \hat{w}, \quad (2.4)$$

$$z_1 p_2^+ \hat{w} = p_3^+ \hat{w} - p_2^+ \hat{w}' + w_1' \hat{w}' + p_2^+ w_1 \cdot \hat{w}, \quad (2.5)$$

$$z_1 p_3^+ \hat{w} = p_4^+ \hat{w} - p_3^+ \hat{w}' + p_3^+ w_1 \cdot \hat{w} + p_2^+ w_1 \cdot \hat{w}' + w_1' \cdot p_2^+ \hat{w} \quad (2.6)$$

and, generally,

$$z_1 p_j^+ \hat{w} = p_{j+1}^+ \hat{w} - p_j^+ \hat{w}' + \sum_{l=0}^{j-1} p_{j-l}^+ w_1 \cdot p_l^+ \hat{w}. \quad (2.7)$$

3. New identities.

Let

$$\mathbf{G}_n = \mathbf{G} - \sum_0^{n-1} p_i^+ z_2^{-i}.$$

In other words, $\mathbf{G}_n$ is the operator $\mathbf{G}$ minus a part of its asymptotic in z_2^{-1} . Let

$$\mathbf{H} = -\partial + \mathbf{G}_1 w_1 + \hat{w} \cdot \text{res}_{z_1}.$$

The statement of the lemma of Sect.2 can be presented as

$$(z_1 - z_2) \mathbf{G}_1 \hat{w} = -\mathbf{G}_1 \hat{w}' - \hat{w}' + \mathbf{G}_1 w_1 \cdot \mathbf{G}_1 \hat{w} + \mathbf{G}_1 w_1 \cdot \hat{w}.$$

Taking into account that $\text{res}_{z_1} \mathbf{G}_1 \hat{w} = \mathbf{G}_1 w_1$, we find the following form of the Fay identity,

$$(z_1 - z_2) \mathbf{G}_1 \hat{w} = \mathbf{H}(\mathbf{G}_1 \hat{w}) - \hat{w}'. \quad (3.1)$$

Proposition 1. *The equation*

$$(z_1 - z_2) n^{(k)} = \mathbf{H}(n^{(k)}) - \partial_k \hat{w} \quad (3.2)$$

has a unique solution

$$n^{(k)} = \sum_1^\infty n_j^{(k)} z_2^{-j}, \quad n_j^{(k)} = P_j^{(k)} \hat{w}$$

where $P_j^{(k)}$ are differential operators in ∂_l having a form $\sum f^{\alpha_1, \dots, \alpha_s} p_{\alpha_1}^+ \dots p_{\alpha_s}^+$ being $\alpha_1, \dots, \alpha_s > 0$, $s > 0$. Coefficients f do not depend on z_1 .

In the case $k = 1, 2$, and 3 the explicit expression for $n^{(k)}$ are

$$n^{(1)} = \mathbf{G}_1 \hat{w}, \quad (3.3a)$$

$$n^{(2)} = 2z_2 \mathbf{G}_2 \hat{w} - (\partial - \mathbf{G}_1 w_1) n_1, \quad (3.3b)$$

$$n^{(3)} = 3z_2^2 \mathbf{G}_3 \hat{w} - (\partial - \mathbf{G}_1 w_1) n_2 - z_2 (\partial - \mathbf{G}_1 w_1) \mathbf{G}_2 \hat{w}. \quad (3.3c)$$

Proof. The equation (3.2) is equivalent to a set of recurrence relations

$$n_j^{(k)} = z_1 n_{j-1}^{(k)} + (n_{j-1}^{(k)})' - \sum_1^{j-2} p_m^+ w_1 \cdot n_{j-1-m}^{(k)} - n_{-1, j-1}^{(k)} \hat{w}, \quad n_1^{(k)} = \partial_k \hat{w} \quad (3.4)$$

where $n_{-1, j-1}^{(k)} = \text{res}_{z_1} n_{j-1}^{(k)}$ and the term with a sum is absent when $j - 2 < 1$. One can use induction. For $j = 1$ the statement is true: one can prove a formula

$$\partial_k = k \sum_{\alpha_1 + \dots + \alpha_s = k} \frac{(-1)^{s-1}}{s} p_{\alpha_1}^+ \dots p_{\alpha_s}^+$$

where the sum runs over all decompositions of k into sums of positive (all $\alpha_l > 0$) integers¹.

¹The proof is: $\sum_1^\infty \partial_k z_2^{-k} / k = \ln \exp \sum_1^\infty \partial_k z_2^{-k} / k = \ln \mathbf{G} = \ln(1 + \mathbf{G}_1) = \sum_1^\infty (-1)^{s-1} \mathbf{G}_1^s / s$. Expanding in powers of z_2^{-1} , one arrives at the required formula.

Let the statement be true for $n_{j-1}^{(k)}$. One can eliminate $z_1 n_{j-1}^{(k)}$ from Eq.(3.4) with the help of the Fay identity (2.7). We find that $n_j^{(k)}$ has a needed form, however without guarantee that there are no terms where all the α_l 's vanish. If this were the case, $n_j^{(k)}$ would contain terms of zero degree in z_1^{-1} . On the other hand, expanding (3.4) in powers of z_1^{-1} and taking the zero degree term, one has: $n_j^{(k)}|_{z_1^0} = n_{-1,j-1}^{(k)} - n_{-1,j-1}^{(k)} = 0$. This proves the proposition for all the j 's and k 's. The uniqueness easily follows from the recurrence formula (3.4).

The additional statement (3.3) for $k = 1$ is just the Fay identity (3.1). The others ($k = 2$ and 3) can be obtained by straightforward but cumbersome calculations, see Appendix. $\square$

4. The universal equation. Let Q be a function. We call the equation

$$f' + 1 - z_2 f + Qf - \mathbf{G}_1 w_1 f = 0 \quad (4.1)$$

universal since it plays a fundamental role for all the constrained hierarchies, as we will see below. This equation has exactly one solution in the form of a series $f_Q = \sum_1^\infty f_j z_2^{-j}$. The coefficients f_j are functions. It is easy to see that $f_1 = 1$, $f_2 = Q$, $f_3 = Q' + Q^2 - w_1'$ etc.

5. k -constraint. The k -constraint is $L_-^k = q\partial^{-1}r$ where q and r satisfy differential equations

$$\partial_n q = L_+^n q, \quad \partial_n r = -L_+^{n*} r \quad (5.1)$$

and L_+^{n*} is a formal adjoint operator. It is well-known that this constraint is compatible with the hierarchy equations. The equation for $\hat{w}(t, \partial)$: $\partial_n \hat{w}(t, \partial) = -L_-^n \hat{w}(t, \partial)$ transforms for $n = k$ to

$$\partial_k \hat{w}(t, \partial) = -q\partial^{-1}r\hat{w}(t, \partial). \quad (5.2)$$

Dividing the last equation by q , and multiplying on the left by ∂ , one gets

$$\partial \circ \partial_k(\hat{w}(t, \partial)) - \frac{q'}{q} \partial_k(\hat{w}(t, \partial)) = -qr\hat{w}(t, \partial).$$

The parentheses in the l.h.s. mean that the operator ∂_k acts on $\hat{w}(t, \partial)$, $q' = \partial(q)$. Rewriting this as

$$\partial_k(\hat{w}(t, \partial))\partial + \partial_k(\hat{w}'(t, \partial)) - \frac{q'}{q} \partial_k \hat{w}(t, \partial) = -qr\hat{w}(t, \partial),$$

one obtains for $\hat{w}(t, z_1)$ the following:

$$z_1 \partial_k(\hat{w}(t, z_1)) + \partial_k(\hat{w}'(t, z_1)) - \frac{q'}{q} \partial_k \hat{w}(t, z_1) = -qr\hat{w}(t, z_1).$$

Finally, let $Q = q'/q$ and $R = qr$, then

$$\partial_k(\hat{w}'(t, z_1)) + (z_1 - Q)\partial_k(\hat{w}(t, z_1)) + R\hat{w}(t, z_1) = 0. \quad (5.3)$$

The first term of expansion in powers of z_1^{-1} yields $R = -\partial_k w_1$.

6. Conditions that $\hat{w}$ belongs to a constrained hierarchy. Suppose one wants to find necessary and sufficient conditions that a Baker function $\psi(t, z_1) = \hat{w}(t, z_1) \exp \xi(t, z_1)$

of the KP hierarchy belongs to the k -constrained subhierarchy. (With some abuse of terminology, we also call $\hat{w}$ a Baker function). This problem can be solved easily. If $\hat{w}$ belongs to the constrained hierarchy then Eq.(5.3) holds and

$$\frac{\partial_k \hat{w}' + z_1 \partial_k \hat{w} - \partial_k w_1 \cdot \hat{w}}{\partial_k \hat{w}} = Q, \quad (6.1)$$

and therefore the left-hand side of this equation is independent of z_1 . Conversely, let the expression (6.1) where $\hat{w}$ is a Baker function of the KP hierarchy not depend on z_1 . Then letting $R = -\partial_k w_1$ and solving the equations $Q = q'/q$ and $R = qr$ with respect to q and r , one easily obtains (5.2). It is known (see [2]) that this implies that q and r satisfy Eq.(5.1), i.e., $\hat{w}$ belongs to the k -constrained hierarchy. Thus, the following criterion holds:

Proposition 2. *A Baker function of the KP hierarchy belongs to the k -constrained hierarchy if and only if the expression (6.1) is independent of z_1 .*

The term with z_1 in (6.1) can be eliminated with the aid of (2.4-7). E.g., for $k = 1$ using (2.4) we have

$$\partial_1 \hat{w}' + z_1 \partial_1 \hat{w} - w_1' \hat{w} = \hat{w}'' + z_1 \hat{w}' - w_1' \hat{w}' = p_2^+ \hat{w}$$

and the condition reads: $p_2^+ \hat{w}/\hat{w}'$ must be independent of z_1 . The same also can be expressed as

$$\begin{vmatrix} w_{i_1}' & w_{i_2}' \\ p_2^+ w_{i_1} & p_2^+ w_{i_2} \end{vmatrix} = 0 \quad (6.2)$$

where i_1 and i_2 are arbitrary.

In the case $k = 2$ we have

$$\partial_3 \hat{w} - z_1 \partial_2 \hat{w} = p_3^+ \hat{w} - w_1' \hat{w}' - \partial_2 w_1 \cdot \hat{w}$$

Then the expression (6.1) takes the form, using (2.5),

$$Q = \frac{\partial_2 \hat{w}' + \partial_3 \hat{w} - p_3^+ \hat{w} + w_1' \hat{w}'}{\partial_2 \hat{w}} = \frac{2p_3^+ \hat{w} - p_2^+ \hat{w}' + w_1' \hat{w}'}{\partial_2 \hat{w}}.$$

We also have used here the known explicit expressions for Schur polynomials:

$$p_2^+ = \frac{1}{2}(\partial^2 + \partial_2), \quad p_3^+ = \frac{1}{3}\partial_3 + p_2^+ \partial - \frac{1}{3}\partial^3.$$

Independence of z_1 of the expression Q is equivalent to

$$\begin{vmatrix} \partial_2 w_{i_1} & \partial_2 w_{i_2} \\ (\partial_3 + \partial_2 \partial - p_3^+ + w_1' \partial) w_{i_1} & (\partial_3 + \partial_2 \partial - p_3^+ + w_1' \partial) w_{i_2} \end{vmatrix} = 0$$

or

$$\begin{vmatrix} (2p_2^+ - \partial^2) w_{i_1} & (2p_2^+ - \partial^2) w_{i_2} \\ (2p_3^+ \hat{w} - p_2^+ \hat{w}' + w_1' \hat{w}') w_{i_1} & (2p_3^+ \hat{w} - p_2^+ \hat{w}' + w_1' \hat{w}') w_{i_2} \end{vmatrix} = 0. \quad (6.3)$$

For $k = 3$, as it can be shown, the condition is

$$\begin{vmatrix} P_1^{(3)} w_{i_1} & P_1^{(3)} w_{i_2} \\ P_2^{(3)} w_{i_1} & P_2^{(3)} w_{i_2} \end{vmatrix} = 0 \quad (6.4)$$

where

$$P_1^{(3)} = 3p_3^+ - 3p_2^+ \partial + \partial^3, \quad P_2^{(3)} = 3p_4^+ - 3p_3^+ \partial + p_2^+ \partial^2 + 3w_1' p_2^+ - 2w_1' \partial^2 - w_1'' \partial.$$

All this is simple and straightforward and would not deserve a special discussion if not the following generalization. As it will be shown in the next sections, infinite sequences of determinant identities can be written in each of cases $k = 1$, $k = 2$ and $k = 3$ such that (6.2), (6.3) and (6.4) are but first terms of them. A similar statement is also true for any k .

7. Main results about constrained hierarchies. Here we make a statement and derive some corollaries of it. In the next section we give a proof.

Proposition 3. *If $n^{(k)}$ are those defined in Sect.3 then for a k -constrained hierarchy we have*

$$\frac{n^{(k)}}{\partial_k \hat{w}} = f_Q, \quad k = 1, 2, 3, \dots \quad (7.1)$$

Here f_Q is the solution of the universal equation, Sect.4, and $Q = q'/q$. Hence the left-hand side of the equation (7.1) does not depend on z_1 .

What we have done actually, is that we constructed some expressions having the asymptotic $z_2^{-1} + Qz_2^{-2} + \dots$ with a property: if the first nontrivial term Q does not depend on z_1 then neither does the whole series.

The numerators of these expressions can be expanded in z_2^{-1} :

$$n^{(k)} = \sum_{j=1}^{\infty} P_j^{(k)} \hat{w} \cdot z_2^{-j}$$

where $P_j^{(k)}$ are differential operators. Namely,

$$P_j^{(1)} = p_j^+, \quad P_j^{(2)} = \sum_{l=1}^{j-1} p_l^+ w_1 \cdot p_{j-l}^+ - p_j^+ \partial + 2p_{j+1}^+$$

and

$$\begin{aligned} P_j^{(3)} = & \sum_{\alpha, \beta, \gamma > 0, \alpha + \beta + \gamma = j} p_\alpha^+ w_1 \cdot p_\beta^+ w_1 \cdot p_\gamma^+ - \sum_{\alpha=1}^{j-1} (2p_\alpha^+ w_1 \cdot p_{j-\alpha}^+ \partial + p_\alpha^+ w_1' \cdot p_{j-\alpha}^+) + 3 \sum_1^{j-1} p_\alpha^+ w_1 \cdot p_{j+1-\alpha}^+ \\ & + p_j^+ \partial^2 - 3p_{j+1}^+ \partial + 3p_{j+2}^+. \end{aligned}$$

Corollary 4. *The equalities*

$$P_j^{(k)} w_i = f_j \cdot \partial_k w_i, \quad j \geq 1$$

hold.

Corollary 5. *There are identities*

$$\begin{vmatrix} P_{j_1}^{(k)} w_{i_1} & P_{j_1}^{(k)} w_{i_2} \\ P_{j_2}^{(k)} w_{i_1} & P_{j_2}^{(k)} w_{i_2} \end{vmatrix} = 0$$

where i_1, i_2, j_1 and j_2 are arbitrary ≥ 1 .

The equalities (6.2-4) are the first of these identities, with the least possible j_1 and j_2 .

8. Proof of the proposition 3. First we prove that the left-hand side of Eq.(7.1) satisfies a slightly different equation

$$f' = -1 + z_2 f + \mathbf{G}_1 w_1 \cdot f - Qf - \frac{\partial_k w_1 \cdot \hat{w}}{\partial_k \hat{w}} (f - f_0) \quad (8.1)$$

where f_0 is the limiting value of f when $z_1 \rightarrow \infty$: $f_0 = n_{-1}^{(k)} / \partial_k w_1$, where $n_{-1}^{(k)} = \text{res}_{z_1} n^{(k)}$. If we manage to prove this, then, passing to the limit $z_1 \rightarrow \infty$ in the equation (8.1), we get for f_0 :

$$f'_0 = -1 + z_2 f_0 + \mathbf{G}_1 w_1 f_0 - Qf_0.$$

Subtracting this equation from Eq.(8.1), we obtain for $g = f - f_0$:

$$g' = z_2 g + \mathbf{G}_1 w_1 \cdot g - Qg - \frac{\partial_k w_1 \cdot \hat{w}}{\partial_k \hat{w}} g.$$

This is a homogeneous equation, g is a double series in z_1^{-1} and z_2^{-1} . Let $g = \sum_{j^*}^\infty g_j z_2^{-j}$ where g_{j^*} is the first non-zero coefficient assuming that it exists. Then expanding the equation for g in powers of z_2^{-1} one gets for the coefficient of $z_2^{-j^*+1}$: $g_{j^*} = 0$ in contradiction to the assumption. This means that $g \equiv 0$. Then $f = f_0$, f does not depend on z_1 and satisfies the universal equation, as required. Thus, it suffices to prove that the left-hand side of Eq.(7.1), which will be denoted as f , satisfies Eq.(8.1).

We have $f = n^{(k)} / \partial_k \hat{w}$. What equation for $n^{(k)}$ does follow from Eq.(7.1)? Firstly,

$$f' = \frac{(n^{(k)})'}{\partial_k \hat{w}} - f \cdot \frac{\partial_k \hat{w}'}{\partial_k \hat{w}}.$$

Eliminating $\partial_k \hat{w}'$ with the help of the constraint equation (5.3) one obtains

$$f' = \frac{(n^{(k)})'}{\partial_k \hat{w}} + (z_1 - Q)f - \frac{\partial_k w_1 \cdot \hat{w}}{\partial_k \hat{w}} f.$$

Substituting this expression for f' in (8.1) one gets the equation for numerators

$$(z_1 - z_2)n^{(k)} = -(n^{(k)})' + \mathbf{G}_1 w_1 \cdot n_k + n_{-1}^{(k)} \hat{w} - \partial_k \hat{w}. \quad (8.2)$$

Here $n_{-1}^{(k)} = f_0 \cdot \partial_k w_1 = \text{res}_{z_1} n^{(k)}$. This equation is nothing less than the equation of the proposition 1, and $n^{(k)}$'s are its solutions. $\square$

Acknowledgement. We are thankful to the *Mathematisches Forschungsinstitut Oberwolfach* whose *RiP* program helped us to accomplish this work.

Appendix. Proof of the Proposition 1 for $k = 2$ and 3 .

Recall the definition of the operator $\mathbf{H}$:

$$\mathbf{H}(v) = -(\partial - \mathbf{G}_1 w_1)v + v_{-1}\hat{w} = -v' + \mathbf{G}_1 w_1 \cdot v + v_{-1}\hat{w},$$

here

$$v_{-1} = \text{res}_{z_1}(v).$$

Thus, for instance,

$$\begin{aligned} n_{-1}^{(1)} &= \mathbf{G}_1 w_1, \\ n_{-1}^{(2)} &= 2z_2 \mathbf{G}_2 w_1 - (\partial - \mathbf{G}_1 w_1)n_{-1}^{(1)}, \\ n_{-1}^{(3)} &= 3z_2^2 \mathbf{G}_3 w_1 - (\partial - \mathbf{G}_1 w_1)n_{-1}^{(2)} - z_2(\partial - \mathbf{G}_1 w_1)\mathbf{G}_2 w_1, \end{aligned}$$

We prepare a few properties of $\mathbf{H}$. For any function v

$$\mathbf{H}(\mathbf{G}_1 w_1 \cdot v) = \mathbf{G}_1 w_1 \cdot \mathbf{H}(v) - \mathbf{G}_1 w_1' \cdot v$$

and

$$-(\partial - \mathbf{G}_1 w_1)\mathbf{H}(v) = \mathbf{H}(-(\partial - \mathbf{G}_1 w_1)v) - v_{-1}\hat{w}'.$$

The following lemmata will be frequently needed.

Lemma 1. *Let*

$$(z_1 - z_2)v = \mathbf{H}(v) - u$$

then

$$(z_1 - z_2)(-(\partial - \mathbf{G}_1 w_1)v) = \mathbf{H}(-(\partial - \mathbf{G}_1 w_1)v) - v_{-1}\hat{w}' + (\partial - \mathbf{G}_1 w_1)u.$$

This is obvious.

Lemma 2. *The following holds:*

$$(z_1 - z_2)\mathbf{G}_1 \hat{w} = \mathbf{H}(\mathbf{G}_1 \hat{w}) - p_1^+ \hat{w}, \quad (1)$$

$$(z_1 - z_2)z_2 \mathbf{G}_2 \hat{w} = \mathbf{H}(z_2 \mathbf{G}_2 \hat{w}) + (-p_2^+ \hat{w} + \mathbf{G}_1 w_1 \cdot p_1^+ \hat{w}), \quad (2)$$

$$\begin{aligned} (z_1 - z_2)z_2^2 \mathbf{G}_3 \hat{w} &= \mathbf{H}(z_2^2 \mathbf{G}_3 \hat{w}) + z_2(\mathbf{G}_1 w_1 \cdot p_1^+ \hat{w}) \\ &\quad + (-p_3^+ \hat{w} + \mathbf{G}_1 w_1 \cdot p_2^+ \hat{w} - w_1' p_1^+ \hat{w}), \end{aligned} \quad (3)$$

Proof. One must replace $\mathbf{G}_2 \hat{w}$ by $\mathbf{G}_1 \hat{w} - z_2^{-1} \hat{w}'$ and $\mathbf{G}_3 \hat{w}$ by $\mathbf{G}_1 \hat{w} - z_2^{-1} \hat{w}' - z_2^{-2} p_2^+ \hat{w}$, then apply equations (3.1) and (2.4-6). Finally, one must go back to $\mathbf{G}_2 \hat{w}$ and $\mathbf{G}_3 \hat{w}$. The calculations are not difficult. $\square$

Now we are proving the proposition. In the case $k = 1$ there is nothing to prove, the required equation coincides with the Fay identity (3.1).

In the case $k = 2$ Lemma 2 yields

$$(z_1 - z_2)2z_2\mathbf{G}_2\hat{w} = 2\mathbf{H}(z_2\mathbf{G}_2\hat{w}) + 2(-p_2^+\hat{w} + 2\mathbf{G}_1w_1 \cdot p_1^+\hat{w}).$$

Lemma 1, starting from (3.1), implies

$$\begin{aligned} (z_1 - z_2)(-(\partial - \mathbf{G}_1w_1)n^{(1)}) &= \mathbf{H}(-(\partial - \mathbf{G}_1w_1)n^{(1)}) \\ &\quad - n_{-1}^{(1)}\hat{w}' + (\partial - \mathbf{G}_1w_1)\hat{w}' \\ &= \mathbf{H}(-(\partial - \mathbf{G}_1w_1)n^{(1)}) - 2\mathbf{G}_1w_1\hat{w}' + \hat{w}''. \end{aligned}$$

A sum of two last equations is exactly what was to prove.

Now, let $k = 3$. Applying Lemma 2 again, we have

$$\begin{aligned} (z_1 - z_2)3z_2^2\mathbf{G}_3\hat{w} &= \mathbf{H}(3z_2^2\mathbf{G}_3\hat{w}) + 3z_2(\mathbf{G}_1w_1 \cdot p_1^+\hat{w}) \\ &\quad + 3(-p_3^+\hat{w} + \mathbf{G}_1w_1 \cdot p_2^+\hat{w} - w_1'p_1^+\hat{w}), \end{aligned}$$

while Lemma 1 and the equation we have just proven for $k = 2$ yield:

$$\begin{aligned} (z_1 - z_2)(-(\partial - \mathbf{G}_1w_1)n^{(2)}) &= \mathbf{H}(-(\partial - \mathbf{G}_1w_1)n^{(2)}) \\ &\quad - n_{-1}^{(2)}\hat{w}' + (\partial - \mathbf{G}_1w_1)\partial_2\hat{w}. \end{aligned}$$

Finally we use the equation for $\mathbf{G}_2\hat{w}$ and apply Lemma 1 to it:

$$\begin{aligned} (z_1 - z_2)z_2(-(\partial - \mathbf{G}_1w_1)\mathbf{G}_2\hat{w}) &= H(z_2(-(\partial - \mathbf{G}_1w_1)\mathbf{G}_2\hat{w})) \\ &\quad - z_2\mathbf{G}_2w_1 \cdot \hat{w}' \\ &\quad + (\partial - \mathbf{G}_1w_1)(p_2^+\hat{w} - \mathbf{G}_1w_1 \cdot \hat{w}'). \end{aligned}$$

On addition of the last three equations we obtain:

$$\begin{aligned} (z_1 - z_2)n^{(3)} &= \mathbf{H}(n^{(3)}) \\ &\quad - 3w_1'\hat{w}' + 3z_2(\mathbf{G}_1w_1 \cdot \hat{w}' - \mathbf{G}_2w_1 \cdot \hat{w}') \\ &\quad - 3p_3^+\hat{w} + \partial\partial_2\hat{w} + \partial p_2^+\hat{w} \\ &\quad + 2\mathbf{G}_1w_1 \cdot p_2^+\hat{w} - \mathbf{G}_1w_1 \cdot \partial_2\hat{w} - \mathbf{G}_1w_1 \cdot \hat{w}'', \end{aligned}$$

which gives the claim by using obvious identities for the Schur polynomials p_2 and p_3 and the equality $\mathbf{G}_1w_1 - \mathbf{G}_2w_1 = z_2^{-1}p_1^+w_1$.

References.

1. Y. Cheng, W. Strampp, B. Zhang, *CMP*, **168** (1995), 117-135.
2. H. Aratyn, Integrable Lax hierarchies, their symmetry reductions and multi-matrix models, Preprint, hep-th 9503211, 1995
3. P. Van Moerbeke, Integrable foundations of string theory, in: *Lectures on Integrable Systems* in Memory of Jean-Louis Verdier, eds. Babilon, Cartier and Kosmann-Schwarzbach, *World Scientific* (1994) 163-268.